

# Please proofread this information

The following is a LaTeX to HTML translation of the abstract information you entered for the Division for Planetary Sciences Meeting. This partial translation is how your abstract will appear online. The LaTeX in your abstract will be fully translated in *The Bulletin of the American Astronomical Society (BAAS)*.

Please proof it and press the COMMIT button on the bottom of this form.

---

## Evolution of Loki's IR Spectrum During its 1997 Eruption from Combined NIMS and Satellite Occultation Radiometry

*J.D. Goguen, A.G. Davies (JPL), P.D. Nicholson, J.E. Van Cleve (Cornell U.), T.L. Hayward (Gemini Obs.), G.J. Veeder, D.L. Blaney, D.L. Matson, T.V. Johnson (JPL), D.W. Toomey (NASA IRTF)*

We use a unique data set consisting of a combination of IR measurements from Palomar, IRTF, and Galileo NIMS to define and model the evolution of Loki's 3 to 11  $\mu\text{m}$  thermal emission spectrum throughout its 1997 eruption. A series of occultations of Io by either Europa or Ganymede on 3/28, 4/11, 4/18, 6/21, 6/28, 7/5, 7/26, 8/17 and 12/10/1997 UT allow isolation of the emission from Loki in ground-based observations. SpectroCAM-10 spectra from Palomar include the critical 8-11  $\mu\text{m}$  spectral region covering the peak of Loki's thermal emission. Galileo NIMS observations include Loki spectra taken during E6 (2/21), G7 (4/4) and C9 (6/28/97) that define the spectrum for wavelengths shorter than 5.2  $\mu\text{m}$ . The 1997 Loki eruption is the optimum data set for this effort due to its occurrence during the Galileo Prime Mission and simultaneously during the series of satellite occultations. It is among the best observed in the number of observations available and with mid-IR spectral coverage resolving the contribution from individual hot spots. This eruption started between 2/21, when NIMS observed low 2.95  $\mu\text{m}$  emission at Loki, and 3/12, when J. Spencer's monitoring program at IRTF reported strong 3.5  $\mu\text{m}$  brightening. The NIMS observations in C9 show that Loki's 3.5  $\mu\text{m}$  flux had faded by 6/28 to about 70% of its value during G7 on 4/4. The well-constrained starting time and the inclusion of 8-11  $\mu\text{m}$  observations spanning 8 months make this data set ideal for applying eruption models that predict the evolution of the spectrum to determine physical parameters of Io's dominant hot spot. The results of these models may have important implications for understanding the close-up Loki observations with NIMS and PPR in the targeted Io encounters during orbits 124-31. This research was funded by grants from the NASA Planetary Astronomy and Jupiter System Data Analysis Programs.

**Presentation Type:** cspo

**Category:** 12. Io and Europa: Galileo results

**Submitter:** Jay Goguen

**Member ID:** 11099

**Presenter email address:** jdg@scn1.jpl.nasa.gov

**Presenter phone:** 818 354 8748

**Correspondent address:** Jet Propulsion Lab MS 183-501 4800 Oak Grove Dr. Pasadena, CA 91109

**SpecialInstructions:** I would be willing to chair either an Io or Galilean Satellite session or a comet session.

**Audio/visual needs:** none

**Member Type:** DPS

**Session chair:** yes

---

If the information is correct, press the "Commit" button below. If the information is *not* correct, please use the **Back** button on your browser to return to the input form and correct the problem.

When you are satisfied with your submission, please print this page for future reference.

Commit

Thank you.